# HIGH SEQUENCE COVERAGE PEPTIDE MAPPING OF A MONOCLONAL ANTIBODY WITH UPLC/MS<sup>E</sup>

Hongwei Xie, Martin Gilar, and John C. Gebler Waters Corporation, Milford, MA, U.S.A.

#### INTRODUCTION

Monoclonal antibodies are a class of recombinant proteins. They are subject to a variety of enzymatic or chemical modifications during expression, purification, and long-term storage. Liquid chromatography (LC)-based peptide mapping is the standard approach to establish protein identity, to identify post-translational modifications (PTMs), and to analyze potential impurities.

Peptide mapping with UltraPerformance LC<sup>®</sup> (UPLC<sup>®</sup>) Technology coupled with data-independent acquisition tandem mass spectrometry (UPLC/MS<sup>E</sup>) has been used to obtain high sequence coverage, and to identify PTMs and variants of enolase.<sup>1-2</sup> The combination of MS<sup>E</sup> for unbiased identification<sup>3</sup> along with the resolution and sensitivity of UPLC<sup>4-5</sup> overcomes difficulties in protein characterization encountered with traditional LC/MS peptide mapping methods and data-dependent acquisition (DDA) LC/MS/MS sequencing tools, including co-elutions, analytical artifacts, and confusion of isobaric or nearly-isobaric peptides. MS<sup>E</sup> has the advantage over DDA in that it does not require peak detection or prior knowledge of the precursor ions.

In this application note, we demonstrate the use of UPLC/MS<sup>E</sup> to characterize a monoclonal antibody digest. The method provides high sequence coverage for both heavy and light chains of the antibody, and identifies modifications in a single analysis. MS<sup>E</sup> spectra were used to identify peptide sequences. The unexpected peptides resulting from non-specific digestion were also identified.

#### EXPERIMENTAL

#### Sample and materials

Commercial monoclonal antibody was in solution (21  $\mu$ g/ $\mu$ L). Iodoacetamide (IAM), dithiothreitol (DTT) and ammonium bicarbonate (NH<sub>4</sub>HCO<sub>3</sub>) were purchased from Sigma Chemical Co. (St. Louis, Mo., U.S.). Sequence-grade trypsin was from Promega Corp. (Madison, Wisc., U.S.), Formic acid (FA) was from EM Science (Gibbstown, N.J., U.S.) and Optima-grade acetonitrile (ACN) was from Fisher Scientific (Pittsburg, Penn., U.S.). RapiGest<sup>™</sup> SF was from Waters Corp. (Milford, Mass., U.S.). The water used in all procedures was from a Millipore Milli-Q purification system (Bedford, Mass., U.S.).

THE SCIENCE

#### Preparation of protein digests

The monoclonal antibody solution was diluted to 5  $\mu$ g/ $\mu$ L with 100 mM NH<sub>4</sub>HCO<sub>3</sub> containing 0.1% RapiGest SF. Proteins in 50  $\mu$ L freshly-prepared solution were digested by adding 5  $\mu$ g of sequencing-grade trypsin and incubating at 37 °C overnight. Before the digestion, 5  $\mu$ L 100 mM DTT was added to reduce disulfide bonds at 60 °C for 30 min. Free cysteine residues were alkylated with 5  $\mu$ L of 200 mM IAM at room temperature for 30 min in the dark. Formic acid (0.5% v/v) was used to quench enzymatic reaction and degrade RapiGest SF. After adding ACN (to 30% v/v), the digest was diluted to a 1.5 pmol/ $\mu$ L final concentration with 5:95 ACN/ water containing 0.1% FA and analyzed by UPLC/MS<sup>E</sup>.

#### UPLC/MS<sup>E</sup> experiments

All analyses were performed using a Waters<sup>®</sup> SYNAPT<sup>™</sup> MS system controlled by MassLynx<sup>™</sup> 4.1 Software. An ACQUITY UPLC<sup>®</sup> System equipped with 2.1 x 150 mm BEH300Å 1.7 µm Peptide Separation Technology C<sub>18</sub> Column was used for the separation. Peptides were eluted with a 90 min gradient (0 to 50 %B). Mobile phase A was 0.1% FA in water, B was 0.1% FA in ACN. The flow rate was 0.2 mL/min, and the column temperature was 40 °C. An auxiliary pump delivered a lockmass solution (100 fmol/µL (GLu1)-fibrinopeptide B (GFP) in 50:50 ACN/water containing 0.1% FA) for mass accuracy reference.

The instrument was operated in the positive ion V-mode. An alternating low collision energy (5 V) and elevated collision energy (ramping from 20 to 40 V) acquisition was used to acquire peptide precursor (MS) and fragmentation (MS<sup>E</sup>) data. Scan time was 0.5 sec (1 sec total duty cycle). The capillary voltage was 3.0 kV, source temperature 100 °C, cone voltage 37 V, cone gas flow 10 L/h. Sampling of the lock spray channel was performed every 1 min.

#### Data processing

The acquired data were processed with Identity<sup>E</sup> Software of ProteinLynx Global SERVER<sup>™</sup> 2.3. The processed data were first searched against a database consisting of monoclonal antibody light and heavy chain sequences, human, rat and mouse proteomes, with trypsin specificity and one potential miscleavage. Then, the data were searched again against the light and heavy chain sequences, without enzyme specificity. Cysteine (C) carbamidomethylation, asparagine (N) deamidation, and methionine (M) oxidation were allowed as optional modifications in these searches.

#### RESULTS AND DISCUSSION

Figure 1 shows the UPLC/MS<sup>E</sup> chromatogram of monoclonal antibody digest (30 pmol injected on-column). The chromatogram features more than 100 detected peaks.

In order to investigate potential protein contaminants in the antibody, the data were searched against a database consisting of human, rat, and mouse proteomes, plus the light and heavy chain sequences. Only the light and heavy chains of the antibody were identified, demonstrating that the sample was not contaminated with other proteins. It has been shown previously that UPLC/MS<sup>E</sup> can identify protein contaminants at levels above 0.1 to 1%.<sup>1</sup>

The tryptic peptides identified are listed in Table 1 and assigned in Figure 1. They comprise 97% sequence of heavy chain and 97.2% sequence of light chain.



Figure 1. Chromatograms of monoclonal antibody tryptic digest. Tryptic peptides originating from heavy (HTxx) and light chain (LTxx) are annotated in blue. Partially tryptic (HPxx or LPxx) and non-tryptic peptides (HNxx or LNxx) are labeled in red. The asterisk indicates peptide with modification. For details, see Tables 1 and 2. Note that some peptides eluted as more than one peak. This is due to separation of glycosilation and deamidation peptide isoforms.

Peptide HT25 was identified as N-linked glycosylated forms. In addition, nine deamidated N sites (seven in heavy chain and two in light chain, two isoforms each as shown in Table 1) were also identified.

After assignment of the tryptic peptides, unknown peaks remain in the chromatogram. These peaks are partially tryptic and non-tryptic peptides, as identified in a subsequent database search against light and heavy chain sequences without enzyme specificity. The identified partially/non-tryptic peptides are presented in Table 2 and shown in Figure 1.

The second search suggests that the longest tryptic peptide HT15 in the heavy chain (63 amino acids; 6712.3 Da) was fragmented into a series of partially tryptic peptides (HP1, HP2, HP15, HP21, HP23, HP25, HP30, HP31, HP32, HP33, HP34) and non-tryptic peptides

(HN1, HN2, HN3, HN4, HN5). All the MS<sup>E</sup> spectra of peptides listed in Table 2 were manually inspected, confirming the validity of identifications.

Figure 2 shows the MS<sup>E</sup> spectra for partially tryptic peptides HP30 and HP15, products of chymotryptic cleavage of tryptic HT15 peptide. When combined, they cover the entire sequence of the HT15.

Two broad peaks were observed in the chromatogram eluting between 65 and 70 minutes (Figure 1). They are proline-rich peptides LT10 (65.5 min) and HT20 (68.3 min). Their peak shapes could be improved by using higher column temperature. MS<sup>E</sup> allows for identification of peptides without complete resolution because it acquires fragment ion data of multiple peptide precursors in parallel.



Figure 2. MS<sup>E</sup> spectra of two partially-tryptic peptides, which cover the whole sequence of the longest tryptic peptide HT15 in heavy chain when combined together. A) HP30, first part of HT15; B) HP15, second part of HT15.

	Peptide <sup>1</sup>	Sequence <sup>2</sup>	Modification	RT (Min)
Heavy-Chain	HT19	SC#DK		2.06
Heavy-Chain	HT29	VSNK		2.14
Heavy-Chain	HT24	TKPR		2.18
Heavy-Chain	HT16	VDK		2.2
Heavy-Chain	HT18	VEPK		2.2
Heavy-Chain	HT27	EYK		2.2
, Heavy-Chain	HT31	TISK		2.2
Heavy-Chain	HT33	GOPR		2.2
Heavy-Chain	нта	OAPGK		4.5
Honw Chain	LT25	EEMTK		9.62
Heavy-Chain	1135	EEMIK		0.03
Heavy-Chain	HI7	YADSVK		13.88
Heavy-Chain	HT39	LTVDK		16.71
Heavy-Chain	HT25*	EEQYNSTYR	Glycosylation N(300)	18.7
Heavy-Chain	HT25*	EEQYNSTYR	Glycosylation N(300)	18.8
Heavy-Chain	HT6*	IYPTNGYTR	Deamidation N(55)	27.19
Heavy-Chain	HT11	AEDTAVYYC#SR		27.93
Heavy-Chain	HT6*	IYPTNGYTR	Deamidation N(55)	28.92
Heavy-Chain	HT9	FTISADTSK		31.44
Heavy-Chain	HT30	ALPAPIEK		32.95
Heavy-Chain	HT21	DTLMISR		32.99
Heavy-Chain	HT42	SI SI SPG		35.19
Heavy-Chain	нт2	L SC#AASGENIK		30.13
leavy=Chain	112			41.02
Heavy-Chain	115	GLEWVAR		41.02
leavy-Chain	HIZ	LSC#AASGENIK	Deamidation N(28)	41.26
neavy-Chain	HI10			41.3
neavy-Chain	H1'10*	NIAYLQMNSLR	Deamidation N(84)	42.62
leavy-Chain	HT2*	LSC#AASGFNIK	Deamidation N(28)	43
leavy-Chain	HT14	STSGGTAALGC#LVK		43.07
leavy-Chain	HT3	DTYIHWVR		43.24
leavy-Chain	HT10*	NTAYLQMNSLR	Deamidation N(84)	43.84
leavy-Chain	HT36	NQVSLTC#LVK		47.31
leavy-Chain	HT13	GPSVFPLAPSSK		48.08
- leavy-Chain	HT36*	NQVSLTC#LVK	Deamidation N(364)	49.2
leavy-Chain	HT41	WQQGNVFSC#SVMHEALHNHYTQK		49.78
Heavy-Chain	HT36*	NOVSI TC#I VK	Deamidation N(364)	50.22
Heavy-Chain	нт1	EVOLVESGGGLVOPGGSLR	Doumidation (Coli)	50.31
leavy-Chain	UT22			50.00
Heavy-Chain	1123			50.56
Heavy-Chain	HIZZ	TPEVIC#VVDVSHEDPEVK	B	50.84
Heavy-Chain	H123*	FNWYVDGVEVHNAK	Deamidation N(289)	51.3
Heavy-Chain	HT23*	FNWYVDGVEVHNAK	Deamidation N(289)	51.9
Heavy-Chain	HT37*	GFYPSDIAVEWESNGQPENNYK	Deamidation N(387)	59.46
leavy-Chain	HT37	GFYPSDIAVEWESNGQPENNYK		59.87
Heavy-Chain	HT37*	GFYPSDIAVEWESNGQPENNYK	Deamidation N(387)	60.6
Heavy-Chain	HT38	TTPPVLDSDGSFFLYSK		62.97
Heavy-Chain	HT20	THTC#PPC#PAPELLGGPSVFLFPPKPK		68.26
Heavy-Chain	HT26	VVSVLTVLHQDWLNGK		71.24
Heavy-Chain	HT26*	VVSVLTVLHQDWLNGK	Deamidation N(318)	71.9
-leavy-Chain	HT35-36	FEMTKNOVSI TC#I VK		72.1
Heavy-Chain	HT26*	VVSVI TVI HODWI NGK	Deemidation N(318)	72.3
Jonay Chain	UT12	WGGDGEYAMDYWGOGTI VTVSSASTK	Deanidation N(010)	72.0
leavy=Chain	1112	NGGEGETAMETWGQGTEVTV33A3TK		00.00
leavy-Chain	HI15	DYFPEPVIVSWNSGALISGVHIFPAVLQS	iGLY	82.28
		SLSSVVTVPSSSLGTQTYIC#NVNHKPSNTK		
ight-Chain	LT20	GEC#		2
ight-Chain	LT16	ADYEK		8.66
ight-Chain	LT6	FSGSR		9.12
ight-Chain	LT19	SFNR		10.62
ight-Chain	LT8	VEIK		14.87
ight-Chain	LT13	VQWK		19.99
.ight-Chain	LT2	VTITC#R		20.71
ight-Chain	LT14	VDNALOSGNSOESVTEODSK		27.28
ight-Chain	I T18	VYAC#EVTHOGI SSPVTK		36.6
ight-Chain	LT3*	ASODVNTAVAWYOOKPGK	Deamidation N/301	37 75
ight Chain	1 T3		20amidad0111(00)	40.62
-ignit=CrialN	1 7 2*		Deamidation 1/00	40.00
ight Chain	L13"		Deamidation N(30)	41.00
ight-Chain	L13S	ASQUVNIAVAWYQQK		42.64
<i>light-Chain</i>	LT1	DIQMTQSPSSLSASVGDR		42.97
<i>light-Chain</i> light-Chain light-Chain		ASQDVNTAVAWYQQK	Deamidation N(30)	44.03
<i>ight-Chain</i> ight-Chain ight-Chain <i>ight-Chain</i>	LT3s*	DETVELETT TI EK		50.16
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	<i>LT3s*</i> LT15	DOTTOLOGILILOK		62.46
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	<i>LT3s*</i> LT15 <i>LT11*</i>	SGTASVVC#LLNNFYPR	Deamidation N(137)	02.70
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	<i>LT3s*</i> LT15 <i>LT11*</i> LT10	SGTASVVC#LLNNFYPR TVAAPSVFIFPPSDEQLK	Deamidation N(137)	65.9
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	<i>LT3s*</i> LT15 <i>LT11*</i> LT10 <i>LT11</i> *	SGTASVVC#LLNNFYPR TVAAPSVFIFPPSDEQLK SGTASVVC#LLNNFYPR	Deamidation N(137) Deamidation N(137)	65.9 67.1
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	LT3s* LT15 LT11* LT10 LT11* LT5	SGTASVVC#LLNNFYPR TVAAPSVFIFPPSDEQLK SGTASVVC#LLNNFYPR LIYSASFLYSGVPSR	Deamidation N(137) Deamidation N(137)	65.9 67.1 67.42
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	LT3s* LT15 LT11* LT10 LT11* LT5 LT4_5	SGTASVCHLINNFYPR TVAAPSVFIFPPSDEQLK SGTASVVCHLINNFYPR LLIYSASFLYSGVPSR APKI LYSASFLYSGVPSP	Deamidation N(137) Deamidation N(137)	65.9 67.1 67.42
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	LT3s* LT15 LT11* LT10 LT11* LT5 LT4-5	SGTASVVC#LLNNFYPR TVAAPSVFIEPPSDEOLK SGTASVVC#LLNNFYPR LLIYSASFLYSGVPSR APKLLIYSASFLYSGVPSR SGTASUACH LINTSZD	Deamidation N(137) Deamidation N(137)	62.76 65.9 67.1 67.42 68.59 72.1
Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain Light-Chain	LT3s* LT15 LT11* LT10 LT11* LT5 LT4-5 LT11	SGTASVVC#LLNNFYPR TVAAPSVFIEPPSDEOLK SGTASVVC#LLNNFYPR LLIYSASFLYSGVPSR APKLLIYSASFLYSGVPSR SGTASVVC#LLNNFYPR SGTASVVC#LLNNFYPR	Deamidation N(137) Deamidation N(137)	62.76 65.9 67.1 67.42 68.59 72.1

Heavy Chain	HP1	IC#NVNHKPSNTK	202	213	Partial	12.70			
Heavy Chain	HP2	HKPSNTK	207	213	Partial	13.88			
Heavy Chain	HP3	STSGGTAALGC#	137	147	Partial	22.06			
Heavy Chain	HP4	EPQVY	348	352	Partial	24.78			
Heavy Chain	HP5	MHEALHNHYTQK	431	442	Partial	25.32			
Heavy Chain	HP6	SVMHEALHNHYTQK	429	442	Partial	29.19			
Heavy Chain	HP7	SGFNIK	25	30	Partial	30.19			
Heavy Chain	HP8	AEDTAVYY	88	95	Partial	33.43			
Heavy Chain	HP9	LSC#AASGF	20	27	Partial	33.65			
Heavy Chain	HP10	NTAYLQMN	77	84	Partial	36.34			
Heavy Chain	HP11	SC#SVMHEALHNHYTQK	427	442	Partial	39.73			
Heavy Chain	HP12	WQQGNVFSC#	420	428	Partial	41.91			
Heavy Chain	HP13	WQQGNVF	420	426	Partial	44.16			
Heavy Chain	HP14	DYWGQGTLVTVSSASTK	108	124	Partial	51.12			
Heavy Chain	HP15	SLSSVVTVPSSSLGTQTYIC#NVNHKPSNTK	184	213	Partial	51.63			
Heavy Chain	HP16	GPSVFPLAPS	125	134	Partial	55.07			
Heavy Chain	HP17	WGGDGFYAM	99	107	Partial	55.41			
Heavy Chain	HP18	FNWYVDGVEVHN	278	289	Partial	55.48			
Heavy Chain	HP19	FNWYVDGVEVH	278	288	Partial	56.32			
Heavy Chain	HP20	YAMDYWGQGTLVTVSSASTK	105	124	Partial	58.38			
Heavy Chain	HP21	DYFPEPVTVSWN	151	162	Partial	62.52			
Heavy Chain	HP22	THTC#PPC#PAPELLGGPSVF	226	244	Partial	62.61			
Heavy Chain	HP23	DYFPEPVTVSWNSGALTSGVH	151	171	Partial	65.33			
Heavy Chain	HP24	PELLGGPSVFLFPPKPK	235	251	Partial	68.30			
Heavy Chain	HP25	DYFPEPVTVSWNSGAL	151	166	Partial	69.25			
Heavy Chain	HP26	TTPPVLDSDGSFFLY	396	410	Partial	71.41			
Heavy Chain	HP27	SVLTVLHQDWLNGK	307	320	Partial	72.29			
Heavy Chain	HP28	VVSVLTVLHQDWLN	305	318	Partial	75.98			
Heavy Chain	HP29	VVSVLTVLHQDWL	305	317	Partial	80.06			
Heavy Chain	HP30	DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY	151	183	Partial	81.36			
Heavy Chain	HP31	DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS	151	186	Partial	82.89			
Heavy Chain	HP32	DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL	151	185	Partial	85.12			
Heavy Chain	HP33	DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYIC#NVN	151	206	Partial	87.55			
Heavy Chain	HP34	DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY	151	201	Partial	87.86			
Heavy Chain	HN1	TVPSSSLGTQTY	190	201	Non	25.15			
Heavy Chain	HN2	SLSSVVTVPSSSLGTQTY	184	201	Non	54.81			
Heavy Chain	HN3	SGALTSGVHTFPAVLQSSGLY	163	183	Non	65.55			
Heavy Chain	HN4	TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY	167	201	Non	77.20			
Heavy Chain	HN5	SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY	163	201	Non	79.24			
Light Chain	LP1	SQESVTEQDSK	159	169	Partial	13.90			
Light Chain	LP2	SASVGDR	12	28	Partial	23.34			
Light Chain	LP3	PPTFGQGTK	95	103	Partial	28.11			
Light Chain	LP4	VYAC#EVTHQGL	191	201	Partial	36.15			
Light Chain	LP5	ASQDVNTAVAWYQQK	25	39	Partial	42.65			
Light Chain	LP6	LLNNFYPR	135	142	Partial	45.40			
Light Chain	LP7	TVAAPSVF	109	116	Partial	46.44			
Light Chain	LP8	SGTDFTLTIS	67	76	Partial	52.68			
Light Chain	LP9	LLIYSASFLYSGVP	170	181	Partial	59.55			
Light Chain	LP10	SGTDFTLTISSLQPEDFATYYC#QQH	67	91	Partial	71.71			
Light Chain	LP11	SGTDFTLTISSLQPEDFATYYC#QQHY	67	92	Partial	73.29			
Light Chain	LP12	SGTDFTLTISSLQPEDFATYY	67	87	Partial	76.65			
Light Chain	LN1	TAVAWYQQK	31	39	Non	35.42			
Light Chain	LN2	DIQMTQSPSSLSA	1	13	Non	40.68			
Light Chain	LN3	SLQPEDFATYYC#QQHY	77	92	Non	53.84			
1 HP - Partially	tryptic pe	ptide from heavy chain, HN - Non-tryptic peptide from heavy chain,							
LP - Partially tryptic peptide from light chain, LN - Non-tryptic peptide from light chain.									
<sup>2</sup> C# - Carbamidomethyl cysteine.									

Start End Cleavage<sup>3</sup> RT (min)

<sup>3</sup> Partial - Partially tryptic cleavage; Non - Non-tryptic cleavage.

Protein

Peptide<sup>1</sup> Sequence<sup>2</sup>

Table 2. Sequences and retention times of identified partially tryptic and non-tryptic peptides.

Table 1. Tryptic peptides identified in the mAb digest. Monoclonal antibody was digested after a prolonged storage in solution.

#### CONCLUSION

The results of the analysis of a monoclonal antibody, with very high sequence coverage for both the heavy and light chains, illustrate that UPLC/MS<sup>E</sup> is capable of characterizing digests of large, complex proteins such as monoclonal antibodies. The unexpected peptides in the sample resulting from non-specific digestion were also identified with high confidence.

MS<sup>E</sup> enables sampling low-abundant components and acquires indiscriminately MS<sup>E</sup> spectrum. The specific conclusions from this study show that:

- High sequence coverage can be obtained for both heavy and light chains of a monoclonal antibody
- Covalent modifications, for example, N-deamidation and N-linked glycosylation, can be identified
- Peptide identifications are supported by high-quality MS<sup>E</sup> spectra

MS<sup>E</sup> collects precursor and fragment ion data of peptides within a single LC run, which enables the method to sequence the LC-eluted peptides. This ensures identification of unknown protein contaminants and PTMs in a single analysis. Therefore, the method may be used for detailed characterization of such proteins without additional MS/MS analyses and sacrificing the analytical efficiency. The method should increase the analytical efficiency in the characterization of therapeutic proteins.

#### References

- Xie HW, Gilar M, Gebler JC. Characterization of Protein Impurities by Peptide Mapping with UPLC/MSE. Waters Application Note. 2009; 720002809en.
- Chakraborty AB, Berger SJ, Gebler JC. Use of an Integrated MS-Multiplexed MS/MS Data Acquisition for High-coverage Peptide Mapping Studies. Rapid Comm. Mass Spec. 2007; 21: 730.
- Silva JC, Denny R, Dorschel CA, Gorenstein M, Kass IJ, Li G-Z, McKenna T, Nold M, Richardson K, Young P, Geromanos S. Quantitative Proteomic Analysis by Mass Retention Time Pairs. Anal. Chem. 2005; 77: 2187.
- 4. Mazzeo JR, Wheat TE, Gillece-Castro BL, Lu Z. Next Generation Peptide Mapping with UPLC. BioPharm International, January 1, 2006.
- Wheat TE, Lu Z, Gillece-Castro BL, Mazzeo JR. Quantitative Aspects of UPLC Peptide Mapping. Waters Application Note. 2007; 720001839en.

# Waters

Waters, UltraPerformance LC, UPLC, and ACQUITY UPLC are registered trademarks of Waters Corporation. SYNAPT, RapiGest, MassLynx, ProteinLynx Global SERVER, and The Science of What's Possible are trademarks of Waters Corporation. All other trademarks are the property of their respective owners. ©2009 Waters Corporation. Produced in the U.S.A. January 2009. 720002821en AG-PDF Waters Corporation 34 Maple Street Milford, MA 01757 U.S.A. T: 1 508 478 2000 F: 1 508 872 1990 www.waters.com

