

Rapid Development and Optimization of Peptide MRM Assays Using High-Sensitivity Xevo TQ-S MS, Skyline, and Proteomics Discovery Libraries

GOAL

To utilize Skyline SRM/MRM Library Explorer and Waters® Xevo® TQ-S formats to enable intuitive, prompt, and seamless development of multiple SRM/MRM peptide transitions and methods from high-density proteomics LC/MS experiments.

BACKGROUND

Targeted LC/MS/MS SRM/MRM validation has the potential to bridge protein biomarker discovery and validation. Candidate proteins and peptides can be retrieved from one or multiple identification repositories and libraries. Typical bottlenecks observed are the selection of the appropriate proteotypic peptides for SRM/MRM quantitation and the development of the associated LC/MS methods for triple quadrupole mass spectrometers. Software plays a vital role in selecting the correct peptides, especially in the case of complex experiments, to measure as a substitute for the target protein. This transitions into measurement of target peptides, where specific optimizations for each are typically performed. Skyline Discovery Library Explorer is an open source document editor for creating and analyzing targeted proteomics experiments.¹ In this technology brief we describe the use of Skyline for the peptide and transition selection process from libraries for method creation for a Xevo TQ-S Mass Spectrometer.

Utilize data from cross platform DDA and DIA proteomics discovery experiments to quickly and easily design robust, highly sensitive, and high-throughput validation MRM assays with Xevo TQ-S.




Figure 1. Skyline Discovery Library Explorer enabling the selection of peptides from a wide range of cross-platform experimental search results, including DDA (Mascot, X!Tandem, etc.), DIA (PLGS MS^E) and spectral libraries (PeptideAtlas, NIST, GPM). In contrast to in-silico methods, which predict tryptic peptides and their fragmentation characteristics, this method is based on experimentally-derived data. The point of entry to the experiment is to know which peptide(s) require monitoring for the protein(s) of interest.

THE SOLUTION

A peptide library was created from the search results of a DDA LC/MS/MS discovery experiment and a common protein database search engine that analyzed tryptically digested cytosolic *E.coli* spiked with four protein digest standards. One or more peptides, shown in Figure 1, were selected and automatically associated with their parent proteins for SRM/MRM method creation. Next, Skyline determined and presented the best candidate transitions. The three most abundant product ions from which a Xevo TQ-S method can be automatically built are shown in the upper and bottom panes of Figure 2. In this instance, a scheduled SRM/MRM method was created using the built-in retention time model of Skyline and empirical observations as input/landmarks for retention time prediction.





Figure 2. The top pane overviews the selected peptides and transitions from a library using Skyline SRM/MRM Editor. Observation of the experimentally derived MS spectra can greatly aid the selection of the most dominant and appropriate transitions.

The left pane shows the transitions that were selected and created to build a Xevo TQ-S MS with MassLynx Software method editor with time scheduled measurements.

The results are summarized in Figure 3, which shows the summed SRM/MRM chromatograms of two transitions per peptide. Also shown are the responses of two background, confirmatory *E. coli* peptides and those of two of the target components. In this example, peptides from yeast alcohol dehydrogenase (ADH) and enolase, were spiked and detected with confident signal-to-noise levels at the 1.25 attomole level on-column.

SUMMARY

The SRM/MRM method development of target standard peptides and background *E. coli* peptides for Xevo TQ-S has been demonstrated using Skyline and discovery search results, libraries, and repositories.

Reference

1. MacLean *et al.* Bioinformatics. 2010 Apr 1;26(7):966-8.




Figure 3. XIC data from a typical experiment, where peptides from enolase and ADH were measured in a complex background of *E. coli* peptides. The XICs for enolase and ADH are the summation of data from two transitions for each peptide and demonstrates the excellent sensitivity and S/N levels typical for this type of methodology. As comparator, XICs for two *E. coli* background peptides are shown in the two middle traces.

Waters

THE SCIENCE OF WHAT'S POSSIBLE.™



Waters and Xevo are registered trademarks of Waters Corporation. The Science of What's Possible and MassLynx are trademarks of Waters Corporation. All other trademarks are the property of their respective owners.

©2012 Waters Corporation. Produced in the U.S.A.
May 2012 720004345EN LB-PDF

Waters Corporation
34 Maple Street
Milford, MA 01757 U.S.A.
T: 1 508 478 2000
F: 1 508 872 1990
www.waters.com